Name:

Enrolment No:



UNIVERSITY OF PETROLEUM AND ENERGY STUDIES End Semester Examination, December 2021

Course: Elements of Modern Physics

Course Code: PHYS 2009G

Programme: B.Sc (H) Chemistry, Mathematics & Geology

Total pages: 2

Semester: III

Max. Marks: 100 Time: 03 hrs.

Instructions:

• All questions are compulsory (**Q9** and **Q11** have an internal choice)

• Use blank paper as rough work to solve the questions in section-A and write only the correct options (type answers, no upload)

• Scientific calculators can be used for calculations.

SECTION-A

S. No.		Marks	CO
Q1.	The de-Broglie wavelength of virus particle of mass 1.5×10^{-15} kg moving at a speed of 2×10^{-3} m/s is	4	CO2
Q2.	Explain the two drawbacks of Rutherford's atomic model.	4	CO1
Q3.	Select all options that satisfy the properties of wave function ψ (a) the wave function must be single-valued (b) the wave function must be discontinuous (c) the wave function must be continuous (d) the wave function must be differentiable (e) the wave function must be infinite (f) the wave function must be finite valued (g) the wave function must be normalizable	4	CO1
Q4.	Calculate the work function in electron volts of a metal, given that the photoelectric threshold wavelength (a) 6200 Å and (b) 5000 Å (answer upto the second decimal) (use $h = 6.62 \times 10^{-34} \text{ J} - \text{s}$)	4	CO3
Q5.	The half-life of Radon is 3.8 days. After how many days will only one-twentieth of the radon sample be leftover. (answer upto the second decimal) (use $\log_{10} 20 = 1.3010$)	4	CO4

SECTION-B (Question No: 9 has an internal choice)

Q6.	Describe the Davisson and Germer experiment to demonstrate the wave character of electrons.	10	CO1
Q7.	Calculate the following for Hydrogen atom by making use of the given physical constants, (a) the velocity of an electron in the ground state (b) the radius of Bohr's orbit in the ground state (c) time taken by the electron to traverse the first Bohr's orbit and (d) the Rydberg constant $h = 6.6 \times 10^{-34} \text{ J} - \text{s},$ $m_e = 9.1 \times 10^{-31} \text{kg},$ $e = 1.6 \times 10^{-19} \text{ Coulomb},$ $c = 3 \times 10^8 \text{ m/s},$ $\varepsilon_0 = 8.86 \times 10^{-12} \text{ coulomb}^2/\text{Newton} - \text{metre}^2$	10	CO2
Q8.	Describe the liquid drop model of the nucleus.	10	CO4
Q9.	Explain the de-Broglie concept of wave velocity and group velocity. Derive a relation between the wave velocity and group velocity. OR What is the uncertainty principle? By applying the uncertainty principle, explain the non-existence of an electron in the atomic nucleus.	10	CO3
	SECTION-C (Question No: 11 has an internal choice)		
Q10.	(a) Explain the binding energy of the nucleus.	10	CO4
	Find the binding energy of an α -particle from the below-given data. Mass of Helium nucleus = 4.001265 a. m. u Mass of proton = 1.007277 a. m. u Mass of neutron = 1.008666 a. m. u 1 a. m. u = 931.4812 MeV	10	CO4
	Find the binding energy of an α -particle from the below-given data. Mass of Helium nucleus = 4.001265 a. m. u Mass of proton = 1.007277 a. m. u Mass of neutron = 1.008666 a. m. u	10	CO4
Q11.	Find the binding energy of an α-particle from the below-given data. Mass of Helium nucleus = 4.001265 a. m. u Mass of proton = 1.007277 a. m. u Mass of neutron = 1.008666 a. m. u 1 a. m. u = 931.4812 MeV (b) Define the range of the α-particle. Explain Gamow's theory of alpha decay with		
Q11.	Find the binding energy of an α-particle from the below-given data. Mass of Helium nucleus = 4.001265 a. m. u Mass of proton = 1.007277 a. m. u Mass of neutron = 1.008666 a. m. u 1 a. m. u = 931.4812 MeV (b) Define the range of the α-particle. Explain Gamow's theory of alpha decay with the necessary diagram. Obtain an expression for the transmission coefficient in a rectangular potential barrier. OR (a) Show that the direction of the recoiled electron in Compton's effect is	10	CO4