

Name:

Enrolment No:



UNIVERSITY OF PETROLEUM AND ENERGY STUDIES

End Semester Examination, December 2018

Programme Name: B. Tech CivilE-Sz-Infra

Semester : III

Course Name : Applied Numerical Methods

Time : 03 hrs

Course Code : MATH 3002

Max. Marks : 100

Nos. of page(s) : 03

Instructions: Attempt all questions from **Section A** (each carrying 4 marks); **Section B** (each carrying 10 marks); **Section C** (each carrying 20 marks).

SECTION A
(Attempt all questions)

Q. No.		Marks	CO												
1.	Estimate the missing figures in the following table: <table border="1" style="margin-left: 20px;"> <tr> <td>x</td> <td>1</td> <td>2</td> <td>3</td> <td>4</td> <td>5</td> </tr> <tr> <td>y</td> <td>2</td> <td>-6</td> <td>7</td> <td>-6</td> <td>32</td> </tr> </table>	x	1	2	3	4	5	y	2	-6	7	-6	32	[4]	CO1
x	1	2	3	4	5										
y	2	-6	7	-6	32										
2.	Establish the operator relation $E \equiv e^{hD}$, where E and D denote the Shifting and Differential operators respectively. (h is the step-length).	[4]	CO1												
3.	Let $x_0=1.5$ be the initial approximation of a root of the equation $x^2 + \log_e x - 2 = 0$. Find the first approximate root of the equation using fixed-point iteration method (iteration method).	[4]	CO2												
4.	Decompose the matrix $A = \begin{bmatrix} 4 & 3 \\ 6 & 3 \end{bmatrix}$ in "LU" form by Crout's decomposition method.	[4]	CO2												
5.	Performing two iteration by Picard method, find an approximate value of $y(0.2)$ for the differential equation $\frac{dy}{dx} = x - y$, with the initial condition $y(0) = 1$.	[4]	CO3												

SECTION B

(Q6-Q8 are compulsory and Q9 has internal choice)

6.	Let a function $f(x)$ be known for $(n+1)$ arguments, namely $x_0, x_1, x_2, \dots, x_n$, which are not necessarily equi-spaced, and $y_j = f(x_j)$, for $j=0, 1, 2, \dots, n$ be the corresponding entries. Derive Newton's divided difference interpolation formula to construct an approximate polynomial $\phi(x)$ of degree less than equal to n which satisfies the conditions $\phi(x_j) = f(x_j) = y_j$, for $j=0, 1, 2, \dots, n$.	10	CO1
7.	Let x_0 be an initial approximation of the root of the equation $f(x) = 0$, where $f(x)$ is	10	CO2

	<p>differentiable at $x=x_0$. Derive the first approximate root x_1 by Newton-Raphson method. Hence find the first approximate root of the equation</p> $\log_e x + x - 3 = 0,$ <p>considering $x_0=2$.</p>		
8.	<p>Use fourth order Runge-Kutta method to solve for $y(1.2)$, considering step-length $h=0.1$ for the differential equation</p> $\frac{dy}{dx} = x^2 + y^2,$ <p>with the initial condition $y(1)=1.5$.</p>	10	CO3
9.	<p>Interchange the equations of the following system to obtain a strictly diagonally dominant system. Then apply Gauss-Seidel method to evaluate an approximate solution, taking the initial approximation as $x_1^{(0)}=1, x_2^{(0)}=0, x_3^{(0)}=1$. Perform three iterations.</p> $x_1 + 5x_2 + 3x_3 = 28$ $12x_1 + 3x_2 - 5x_3 = 1$ $3x_1 + 7x_2 + 13x_3 = 76.$ <p style="text-align: center;">OR</p> <p>Using Gauss-Jacobi method, solve the following system of equations starting with initial solution as $x_1^{(0)} = \frac{9}{5}, x_2^{(0)} = \frac{4}{5}, x_3^{(0)} = \frac{6}{5}$. Perform three iterations.</p> $5x_1 - x_2 = 9$ $-x_1 + 5x_2 - x_3 = 4$ $-x_2 + 5x_3 = -6.$	10	CO2
SECTION-C			
(Q10 is compulsory; Q11 has two parts, Q11.A and Q11.B, both have internal choices)			
10.A	<p>Evaluate the integration $\int_0^{0.8} \frac{\sin x}{x} dx$, using (i) Simpson's $\frac{1}{3}$ rule and (ii) Simpson's $\frac{3}{8}$ rule, by dividing the intervals $[0, 0.8]$ into six equal parts.</p>	10	CO1
10.B	<p>Consider a matrix $A = \begin{bmatrix} 4 & 2 & 6 \\ 2 & 82 & 39 \\ 6 & 39 & 26 \end{bmatrix}$.</p> <ol style="list-style-type: none"> Is it possible to find LU-decomposition for the matrix A in form of $A = LL^T$, where L is a lower triangular matrix, by Cholesky method? If yes, decompose the matrix in form of $A = LL^T$. Hence, find the solution of the following system by Cholesky method. $4x_1 + 2x_2 + 6x_3 = 16$ $2x_1 + 82x_2 + 39x_3 = 206$ $6x_1 + 39x_2 + 26x_3 = 113$ 	10	CO2

<p>11.A</p>	<p>Intensity of radiation is directly proportional to the amount of remaining radioactive substance. The differential equation is</p> $\frac{dy}{dx} = -ky, \text{ where } k=0.01.$ <p>Given that $x_0=0$ and $y_0=100$. Determine how much substance will remain at the moment $x=100$, using Modified Euler's method with the step-length $h=50$.</p> <p style="text-align: center;">OR</p> <p>Solve the elliptic equation $u_{xx}+u_{yy}=0$ for the following square mesh with boundary values as shown in the figure by Liebmann's iteration process. Perform five iterations.</p> <div style="text-align: center;"> </div>	<p>10</p>	<p>CO3</p>
<p>11.B</p>	<p>Solve $\frac{\partial u}{\partial t} = \frac{1}{2} \frac{\partial^2 u}{\partial x^2}$ with the conditions $u(0,t)=0, u(4,t)=0, u(x,0)=x(4-x)$ taking $h=1$ and employing Bender-Schmidt method. Continue the solution through five time steps.</p> <p style="text-align: center;">OR</p> <p>Using Crank-Nicholson's method, solve $u_{xx}=16u_t, 0 < x < 1, t > 0$, given that $u(x,0)=0, u(0,t)=0, u(1,t)=50t$. Compute u for two steps in t direction taking $h=\frac{1}{4}$.</p>	<p>10</p>	<p>CO3</p>

Name:	
Enrolment No:	

UNIVERSITY OF PETROLEUM AND ENERGY STUDIES

End Semester Examination, December 2018

Programme Name: B. Tech CivilE-Sz-Infra	Semester : III
Course Name : Applied Numerical Methods	Time : 03 hrs
Course Code : MATH 3002	Max. Marks : 100
Nos. of page(s) : 03	

Instructions: Attempt all questions from **Section A** (each carrying 4 marks); **Section B** (each carrying 10 marks); **Section C** (each carrying 20 marks).

SECTION A
(Attempt all questions)

Q. No.		Marks	CO												
1.	Estimate the missing figures in the following table: <table border="1" style="margin-left: auto; margin-right: auto; border-collapse: collapse;"> <tr> <td style="padding: 2px 10px;">x</td> <td style="padding: 2px 10px;">1</td> <td style="padding: 2px 10px;">2</td> <td style="padding: 2px 10px;">3</td> <td style="padding: 2px 10px;">4</td> <td style="padding: 2px 10px;">5</td> </tr> <tr> <td style="padding: 2px 10px;">y</td> <td style="padding: 2px 10px;">2</td> <td style="padding: 2px 10px;">5</td> <td style="padding: 2px 10px;">7</td> <td style="padding: 2px 10px;">-6</td> <td style="padding: 2px 10px;">32</td> </tr> </table>	x	1	2	3	4	5	y	2	5	7	-6	32	[4]	CO1
x	1	2	3	4	5										
y	2	5	7	-6	32										
2.	Establish the operator relation $\Delta - \nabla \equiv \Delta \nabla$, where Δ and ∇ denote the Newton forward and Backward operators respectively.	[4]	CO1												
3.	Find the value of $(17)^{\frac{1}{3}}$ after four iteration, considering the initial interval as $[2, 3]$ by Bisection method.	[4]	CO2												
4.	Decompose the matrix $A = \begin{bmatrix} 2 & 3 \\ 6 & 3 \end{bmatrix}$ in "LU" form by Crout's decomposition method.	[4]	CO2												
5.	Classify the partial differential equation $a^2 u_{xx} = u_{tt}$, where a is constant.	[4]	CO3												

SECTION B
(Q6-Q8 are compulsory and Q9 has internal choice)

6.	Let a function $f(x)$ be known for $(n+1)$ arguments, namely $x_0, x_1, x_2, \dots, x_n$, which are not necessarily equi-spaced, and $y_j = f(x_j)$, for $j=0, 1, 2, \dots, n$ be the corresponding entries. Derive Lagrange interpolation formula to construct an approximate polynomial $L(x)$ of degree less than equal to n which satisfies the conditions $L(x_j) = f(x_j) = y_j$, for $j=0, 1, 2, \dots, n$.	10	CO1
7.	Let x_0 be an initial approximation of the root of the equation $f(x) = 0$, where $f(x)$ is differentiable at $x = x_0$. Derive the first approximate root x_1 by Newton-Raphson method. Hence find the first approximate root of the equation	10	CO2

	$2x + \log_{10}x - 7 = 0,$ considering $x_0 = 3.5.$		
8.	Intensity of radiation is directly proportional to the amount of remaining radioactive substance. The differential equation is $\frac{dy}{dx} = -ky, \text{ where } k = 0.01.$ Given that $x_0 = 0$ and $y_0 = 100$. Determine how much substance will remain at the moment $x = 100$, using Modified Euler's method with the step-length $h = 50$.	10	CO3
9.	Interchange the equations of the following system to obtain a strictly diagonally dominant system. Then apply Gauss-Jacobi method to evaluate an approximate solution, taking the initial approximation as $x_1^{(0)} = 1, x_2^{(0)} = 0, x_3^{(0)} = 1$. Perform three iterations. $x_1 + 5x_2 + 3x_3 = 28$ $12x_1 + 3x_2 - 5x_3 = 1$ $3x_1 + 7x_2 + 13x_3 = 76.$ <p style="text-align: center;">OR</p> Interchange the equations of the following system to obtain a strictly diagonally dominant system. Then using Gauss-Seidal method, solve the following system of equations starting with initial solution as $x_1^{(0)} = \frac{9}{5}, x_2^{(0)} = \frac{4}{5}, x_3^{(0)} = \frac{6}{5}$. Perform three iterations. $-x_1 + 5x_2 - x_3 = 4$ $5x_1 - x_2 = 9$ $-x_2 + 5x_3 = -6.$	10	CO2
SECTION-C			
(Q10 is compulsory; Q11 has two parts, Q11.A and Q11.B, both have internal choices)			
10.A	Evaluate the integration $\int_0^{\frac{\pi}{2}} \sqrt{1 - k \sin^2 \phi} d\phi, k = 0.162,$ by Simpson's $\frac{1}{3}$ rule, dividing the interval $0 \leq \phi \leq \frac{\pi}{2}$ into six equal subintervals.	10	CO1
10.B	Consider a matrix $A = \begin{bmatrix} 4 & 2 & 6 \\ 2 & 82 & 39 \\ 6 & 39 & 26 \end{bmatrix}.$ <ol style="list-style-type: none"> Is it possible to find LU-decomposition for the matrix A in form of $A = LL^T,$ where L is a lower triangular matrix, by Cholesky method? If yes, decompose the matrix in form of $A = LL^T.$ Hence, find the solution of the following system by Cholesky method. $4x_1 + 2x_2 + 6x_3 = 16$ $2x_1 + 82x_2 + 39x_3 = 206$ $6x_1 + 39x_2 + 26x_3 = 113$ 	10	CO2

11.A	<p>Use fourth order Runge-Kutta method to solve for $y(1.2)$, considering step-length $h=0.1$ for the differential equation</p> $\frac{dy}{dx} = x^2 + y^2,$ <p>with the initial condition $y(1)=1.5$.</p> <p style="text-align: center;">OR</p> <p>Solve the Poisson's equation $u_{xx} + u_{yy} = -10(x^2 + y^2 + 10)$ over the square mesh with sides $x=0, y=0, x=3, y=3$ with $u=0$ on the boundary and mesh length 1. Perform three iterations by Gauss Seidal method to solve the linear equations in u.</p>	10	CO3
11.B	<p>Solve $\frac{\partial u}{\partial t} = \frac{1}{2} \frac{\partial^2 u}{\partial x^2}$ with the conditions $u(0,t)=0, u(4,t)=0, u(x,0)=x(4-x)$ taking $h=1$ and employing Bender-Schmidt method. Continue the solution through five time steps.</p> <p style="text-align: center;">OR</p> <p>Using Crank-Nicholson's method, solve $u_{xx} = 16u_t, 0 < x < 1, t > 0$, given that $u(x,0)=0, u(0,t)=0, u(1,t)=50t$. Compute u for two steps in t direction taking $h = \frac{1}{4}$.</p>	10	CO3

*******GOOD LUCK*******